A Kinetic Framework for Fluids with Ordering

Umberto Zerbinati*, joint work with: J. A. Carillo* , P. E. Farrell*, A. Medaglia*.

*Mathematical Institute – University of Oxford

Bath Royal Literary and Scientific Institution, Internal Seminar, 6th June 2025

Oxford Mathematics



Institute



Partially ordered fluids are ubiquitous in nature and have a wide range of applications.

ORDERED FLUIDS





Partially ordered fluids are ubiquitous in nature and have a wide range of applications.

 Liquid crystals, here depicted in nematic and smectic phases.

ORDERED FLUIDS





Partially ordered fluids are ubiquitous in nature and have a wide range of applications.

- Liquid crystals, here depicted in nematic and smectic phases.
- Ferrofluids, i.e. a colloidal suspension made of nanoscale ferromagnetic or ferrimagnetic particles.



ORDERED FLUIDS





Partially ordered fluids are ubiquitous in nature and have a wide range of applications.

- Liquid crystals, here depicted in nematic and smectic phases.
- Ferrofluids, i.e. a colloidal suspension made of nanoscale ferromagnetic or ferrimagnetic particles.
- Gas saturated magma melts and other fluids with non-diffusive bubbles.



A kinetic theory approach seeks to describe the evolution of the fluid using a statistical description and knowledge of the microscopic interactions.

A KINETIC THEORY APPROACH



- A kinetic theory approach seeks to describe the evolution of the fluid using a statistical description and knowledge of the microscopic interactions.
- Typically, this is done by considering only binary interactions between fluid constituents. This is a good approximation for dilute systems.

J. Am. Chem. Soc. 2011 133 (8), 2346-2349 (A. Kuijk, A. van Blaaderen, A. Imhof). ArXiv First order non-instantaneous corrections in collisional kinetic alignment models, 2025 (L.Kanzler, C. Moschella, C. Schmeiser)



Kinetic Theory of Ordered Fluids

A KINETIC THEORY APPROACH



- A kinetic theory approach seeks to describe the evolution of the fluid using a statistical description and knowledge of the microscopic interactions.
- Typically, this is done by considering only binary interactions between fluid constituents. This is a good approximation for dilute systems.
- In certain cases, the partial ordering of the fluid is a consequence of the dilute nature of the system.

J. Am. Chem. Soc. 2011 133 (8), 2346-2349 (A. Kuijk, A. van Blaaderen, A. Imhof). ArXiv First order non-instantaneous corrections in collisional kinetic alignment models, 2025 (L.Kanzler, C. Moschella, C. Schmeiser)



A KINETIC THEORY APPROACH



- A kinetic theory approach seeks to describe the evolution of the fluid using a statistical description and knowledge of the microscopic interactions.
- Typically, this is done by considering only binary interactions between fluid constituents. This is a good approximation for dilute systems.
- In certain cases, the partial ordering of the fluid is a consequence of the dilute nature of the system.

Onsager's Approach To Liquid Crystals

Onsager explained the emergence of nematic ordering by a truncation of the Mayer cluster expansion, valid for dilute systems. J. Am. Chem. Soc. 2011 133 (8), 2346-2349 (A. Kuijk, A. van Blaaderen, A. Imhof). ArXiv First order non-instantaneous corrections in collisional kinetic alignment models, 2025 (L.Kanzler, C. Moschella, C. Schmeiser)





OXFORD Mathematical nstitute

Order Parameter Manifold

We say that the tuple $(\mathcal{M}, \mathcal{A})$ is an order parameter manifold if \mathcal{M} is a smooth manifold with a fixed parametrization, and \mathcal{A} is a Lie group action of SO(*d*) on \mathcal{M} , i.e. the map \mathcal{A} is smooth enough to be differentiable.

Continua with microstructure, (G. Capriz), Differential geometry and continuum mechanics, (G. Capriz, R.J. Knops).

Order Parameter Manifold

We say that the tuple $(\mathcal{M}, \mathcal{A})$ is an order parameter manifold if \mathcal{M} is a smooth manifold with a fixed parametrization, and \mathcal{A} is a Lie group action of SO(*d*) on \mathcal{M} , i.e. the map \mathcal{A} is smooth enough to be differentiable.

Furthermore, we say that a field $\nu : \mathbb{E}^d \to \mathcal{M}$ is an order parameter field if $\forall \underline{c} \in \mathbb{R}^d$ and $\forall \underline{Q} \in SO(d)$ we have

$$u(\underline{Q}\mathbf{x}+\underline{c})=\mathcal{A}(\underline{Q},\nu(\mathbf{x})),\quad \forall\mathbf{x}\in\mathbb{E}^{d}.$$

Continua with microstructure, (G. Capriz), Differential geometry and continuum mechanics, (G. Capriz, R.J. Knops).



Order Parameter Manifold

We say that the tuple $(\mathcal{M}, \mathcal{A})$ is an order parameter manifold if \mathcal{M} is a smooth manifold with a fixed parametrization, and \mathcal{A} is a Lie group action of SO(d) on \mathcal{M} , i.e. the map \mathcal{A} is smooth enough to be differentiable.

Furthermore, we say that a field $\nu : \mathbb{E}^d \to \mathcal{M}$ is an order parameter field if $\forall \underline{c} \in \mathbb{R}^d$ and $\forall \underline{Q} \in SO(d)$ we have

$$u(\underline{Q}\mathbf{x}+\underline{c})=\mathcal{A}(\underline{Q},\nu(\mathbf{x})),\quad \forall\mathbf{x}\in\mathbb{E}^{d}.$$

Continua with microstructure, (G. Capriz), Differential geometry and continuum mechanics, (G. Capriz, R.J. Knops).

We need to understand what manifold *M* captures the nature of the order parameters.



Order Parameter Manifold

We say that the tuple $(\mathcal{M}, \mathcal{A})$ is an order parameter manifold if \mathcal{M} is a smooth manifold with a fixed parametrization, and \mathcal{A} is a Lie group action of SO(d) on \mathcal{M} , i.e. the map \mathcal{A} is smooth enough to be differentiable.

Furthermore, we say that a field $\nu : \mathbb{E}^d \to \mathcal{M}$ is an order parameter field if $\forall \underline{c} \in \mathbb{R}^d$ and $\forall \underline{Q} \in SO(d)$ we have

$$u(\underline{Q}\mathbf{x}+\underline{c})=\mathcal{A}(\underline{Q},\nu(\mathbf{x})),\quad \forall\mathbf{x}\in\mathbb{E}^{d}.$$

Continua with microstructure, (G. Capriz), Differential geometry and continuum mechanics, (G. Capriz, R.J. Knops).

- We need to understand what manifold *M* captures the nature of the order parameters.
- We need to understand the action of rotations on the manifold *M*.







Variational Theories for Liquid Crystals, (E. Virga), The Physics of Liquid Crystals, (P.G. de Gennes, J. Prost).

We can represent the state of a calamitic molecule using the set of Euler angles θ, φ, ψ.





Variational Theories for Liquid Crystals, (E. Virga), The Physics of Liquid Crystals, (P.G. de Gennes, J. Prost).

- We can represent the state of a calamitic molecule using the set of Euler angles θ, φ, ψ.
- We can also represent the state of a calamitic molecule using a director field <u>v</u> ∈ S².





Variational Theories for Liquid Crystals, (E. Virga), The Physics of Liquid Crystals, (P.G. de Gennes, J. Prost).

- We can represent the state of a calamitic molecule using the set of Euler angles θ, φ, ψ.
- We can also represent the state of a calamitic molecule using a director field <u>ν</u> ∈ S².
- For head-tail symmetric calamitic molecules, we can use ℝP².

Oxford Mathematics



Embedding theorems

Any compact orientable 2-manifold can be embedded in ℝ³. *Curves and Surfaces*, (M. Abate, F. Tovena), *Topology*, (M. Manetti).



Any compact orientable 2-manifold can be embedded in ℝ³. Curves and Surfaces, (M. Abate, F. Tovena), Topology, (M. Manetti).

\$ We can embed the director field $\underline{\nu}$ in \mathbb{R}^3 and work with a vector space structure.



Embedding theorems

- Any compact orientable 2-manifold can be embedded in ℝ³.
- ▶ The real projective space ℝP² can not be embedded in ℝ³.

Curves and Surfaces, (M. Abate, F. Tovena), *Topology*, (M. Manetti).

\$ We can embed the director field $\underline{\nu}$ in \mathbb{R}^3 and work with a vector space structure.



Embedding theorems

- Any compact orientable 2-manifold can be embedded in ℝ³.
- ▶ The real projective space ℝP² can not be embedded in ℝ³.
- ▶ The real projective space ℝP² can be embedded in ℝ⁴.

Curves and Surfaces, (M. Abate, F. Tovena), Topology, (M. Manetti).

Solution We can embed the director field <u>ν</u> in ℝ³ and work with a vector space structure.



Embedding theorems

- Any compact orientable 2-manifold can be embedded in ℝ³.
- ▶ The real projective space \mathbb{RP}^2 can not be embedded in \mathbb{R}^3 .
- ► The real projective space ℝP² can be embedded in ℝ⁴.

Curves and Surfaces, (M. Abate, F. Tovena), *Topology*, (M. Manetti).

- Solution We can embed the director field <u>ν</u> in ℝ³ and work with a vector space structure.
- **\$\$** We can embed the real projective space \mathbb{RP}^2 in \mathbb{R}^4 and work with a vector space structure.



THE MICROSCOPIC WORLD



LAGRANGIAN MECHANICS OF THE CONSTITUENTS



We will here assume that the fluid is composed of a set of constituents, each of which is described by a position \mathbf{x}_i , a velocity $\underline{\nu}_i$, the order parameter ν_i and its total time derivative $\underline{\dot{\nu}}_i$.



LAGRANGIAN MECHANICS OF THE CONSTITUENTS



We will here assume that the fluid is composed of a set of constituents, each of which is described by a position \mathbf{x}_i , a velocity \underline{v}_i , the order parameter ν_i and its total time derivative $\underline{\dot{\nu}}_i$.

$$\mathcal{L}_i \coloneqq \frac{1}{2}m_1(\underline{\dot{x}}_i \cdot \underline{\dot{x}}_i) + \frac{1}{2}\underline{\dot{\nu}}_i \cdot \underline{\Omega}_i(\nu_i)\underline{\dot{\nu}}_i.$$



LAGRANGIAN MECHANICS OF THE CONSTITUENTS

We will here assume that the fluid is composed of a set of constituents, each of which is described by a position \mathbf{x}_i , a velocity v_i , the order parameter v_i and its total time derivative $\underline{\dot{\nu}}_i$.

$$\mathcal{L}_i \coloneqq rac{1}{2} m_1(\dot{\underline{x}}_i \cdot \dot{\underline{x}}_i) + rac{1}{2} \dot{\underline{\nu}}_i \cdot \underline{\Omega}_i(
u_i) \dot{\underline{\nu}}_i.$$

We assume the interaction between the constituents is given by a potential $\mathcal{W}(|\mathbf{x}_i - \mathbf{x}_i|, \nu_i, \nu_i)$, i.e.

$$\mathcal{L}_{i,j} = \mathcal{L}_i(\mathbf{x}_i, \Xi_i) + \mathcal{L}_j(\mathbf{x}_j, \Xi_j) + \mathcal{W}(|\mathbf{x}_i - \mathbf{x}_j|, \nu_i, \nu_j),$$

where $\Xi_i := (\underline{v}_i, \nu_i, \underline{\dot{\nu}}_i)$.





Oxford





If a Lagrangian \mathcal{L} is invariant under a group action with infinitesimal generators G then

$$\frac{d}{dt}\left(\frac{\partial \mathcal{L}}{\partial \dot{q}_{1,2}} \cdot \boldsymbol{G}\right) = 0, \qquad \boldsymbol{q}_{1,2} = \left(\mathbf{x}_1, \mathbf{x}_2, \nu_1, \nu_2\right).$$

In other words for any physical symmetry of the system, there is a conserved quantity.

Analytical Mechanics: An Introduction, (A. Fasano, S. Marmi).





If a Lagrangian \mathcal{L} is invariant under a group action with infinitesimal generators G then

$$\frac{d}{dt}\left(\frac{\partial \mathcal{L}}{\partial \dot{q}_{1,2}} \cdot G\right) = 0, \qquad q_{1,2} = (\mathbf{x}_1, \mathbf{x}_2, \nu_1, \nu_2).$$

In other words for any physical symmetry of the system, there is a conserved quantity.

Analytical Mechanics: An Introduction, (A. Fasano, S. Marmi).

The Lagrangian *L* is invariant under translations, i.e. the linear momentum is conserved.



If a Lagrangian \mathcal{L} is invariant under a group action with infinitesimal generators G then

$$\frac{d}{dt}\left(\frac{\partial \mathcal{L}}{\partial \dot{q}_{1,2}} \cdot G\right) = 0, \qquad q_{1,2} = (\mathbf{x}_1, \mathbf{x}_2, \nu_1, \nu_2).$$

In other words for any physical symmetry of the system, there is a conserved quantity.

Analytical Mechanics: An Introduction, (A. Fasano, S. Marmi).

- The Lagrangian *L* is invariant under translations, i.e. the linear momentum is conserved.
- The Lagrangian L is independent of time and the kinetic energy is a homogeneous quadratic form of the conjugate moments, i.e. the energy is conserved.



Infinitesimal Generator of ${\cal A}$

For fixed $u \in \mathcal{M}$, the orbit map

$$\mathcal{A}_{
u}:\mathsf{SO}(3) o\mathsf{SO}(3)
u,\quad \underline{Q}\mapsto\mathcal{A}(\underline{Q},
u),$$

is differentiable at the identity.

```
We will denote by A_{\nu} : SO(3) \to T_{\nu}\mathcal{M} the differ-
ential of \mathcal{A}_{\nu} at the identity.
Composing the canonical isomorphism \mathbb{R}^3 \to SO(3) with the differential of the orbit map we
obtain a map A_{\nu} : \mathbb{R}^3 \to T_{\nu}\mathcal{M}.
```

Continua with microstructure, (G. Carpiz), Differential geometry and continuum mechanics, (G. Carpiz, R.J. Knops).



Infinitesimal Generator of ${\cal A}$

For fixed $u \in \mathcal{M}$, the orbit map

$$\mathcal{A}_{
u}:\mathsf{SO}(3)
ightarrow\mathsf{SO}(3)
u,\quad \underline{Q}\mapsto\mathcal{A}(\underline{Q},
u),$$

is differentiable at the identity.

We will denote by $A_{\nu} : SO(3) \to T_{\nu}\mathcal{M}$ the differential of \mathcal{A}_{ν} at the identity. Composing the canonical isomorphism $\mathbb{R}^3 \to SO(3)$ with the differential of the orbit map we obtain a map $A_{\nu} : \mathbb{R}^3 \to T_{\nu}\mathcal{M}$. Continua with microstructure, (G. Carpiz), Differential geometry and continuum mechanics, (G. Carpiz, R.J. Knops).

Assuming that the Lagrangian \mathcal{L} is frame-indifferent, i.e. invariant under the action of SO(3), we have:

 $G = (\underline{r} \times \mathbf{x}, \underline{r} \times \underline{x}, A_{\nu}\mathbf{r}, A_{\nu}\underline{r}),$

where \underline{r} is the rotation axis. Thus, the angular momentum is conserved.

AN EXAMPLE: ANGULAR MOMENTUM NEMATIC LIQUID CRYSTALS



For segment like molecules the classical we have $\underline{\Omega}(\underline{\nu}) = I$, where I is the Identity. Thus, Noether's theorem implies the conservation of the following quantity:

$$m_1 \mathbf{x}_1 \times \underline{p}_1 + \nu \times \underline{\dot{\nu}}_1 + m_2 \mathbf{x}_1 \times \underline{p}_2 + \nu \times \underline{\dot{\nu}}_2$$

Let $\underline{\omega}$ be the angular velocity of the segment, using the triple cross product together with the well-known property of segment like rigid bodies that $\underline{\dot{\nu}}_i = \underline{\omega} \times \underline{\nu}_i$ we can rewrite one term of the previous expression as

$$\underline{\nu}_i \times \underline{\dot{\nu}_i} = \underline{\nu}_i \times \underline{\omega}_i \times \underline{\nu}_i = (\underline{\nu}_i \cdot \underline{\nu}_i)\underline{\omega} - (\underline{\nu}_i \cdot \underline{\omega}_i)\underline{\nu}_i = \underline{\omega}_i - (\underline{\nu}_i \cdot \underline{\omega}_i)\underline{\nu}_i = \mathbb{I}_i\underline{\omega},$$

where used the fact that the inertia tensor of a segment is $\mathbb{I}_i := I - \underline{\nu}_i \otimes \underline{\nu}_i$. Therefore, we retrieved the classical definition of angular momentum, i.e.

$$\mathbf{x}_1 \times \underline{p}_1 + \mathbb{I}_1 \underline{\omega}_1 + \mathbf{x}_2 \times \underline{p}_2 + \mathbb{I}_2 \underline{\omega}_2,$$

BBGKY HIERARCHY



HAMILTONIAN MECHANICS OF THE CONSTITUENTS



We introduce the Hamiltonian formalism associated to the Lagrangian \mathcal{L} introduced in the previous section. As usual, we introduce the conjugate momenta to the generalised coordinates, i.e.

$$\underline{p_i} \coloneqq \frac{\partial \mathcal{L}}{\partial \dot{\mathbf{x}}_i} = m \dot{\mathbf{x}}_i, \qquad \underline{s_i} \coloneqq \frac{\partial \mathcal{L}}{\partial \underline{\dot{\nu}}_i} = \underline{\underline{\Omega}}(\nu) \, \underline{\dot{\nu}}_i.$$

We then introduce the Hamiltonian \mathcal{H} of the full system of N constituents, only interacting in pairs, as

$$\mathcal{H} \coloneqq \sum_{i=1}^{N} \frac{1}{2m} \underline{p_i} \cdot \underline{p_i} + \frac{1}{2} \underline{\varsigma_i} \cdot \underline{\Omega}(\nu)^{-1} \underline{\varsigma_i} + \sum_{1 \leq i < j \leq N} \mathcal{W}(|\mathbf{x}_i - \mathbf{x}_j|, \nu_i, \nu_j).$$



We introduce the Hamiltonian formalism associated to the Lagrangian \mathcal{L} introduced in the previous section. As usual, we introduce the conjugate momenta to the generalised coordinates, i.e.

$$\underline{p_i} \coloneqq \frac{\partial \mathcal{L}}{\partial \dot{\mathbf{x}}_i} = m \dot{\mathbf{x}}_i, \qquad \underline{s_i} \coloneqq \frac{\partial \mathcal{L}}{\partial \underline{\dot{\nu}}_i} = \underline{\underline{\Omega}}(\nu) \, \underline{\dot{\nu}}_i.$$

We then introduce the Hamiltonian \mathcal{H} of the full system of N constituents, only interacting in pairs, as

$$\mathcal{H} \coloneqq \sum_{i=1}^{N} \frac{1}{2m} \underline{p_i} \cdot \underline{p_i} + \frac{1}{2} \underline{\varsigma_i} \cdot \underline{\Omega}(\nu)^{-1} \underline{\varsigma_i} + \sum_{1 \leq i < j \leq N} \mathcal{W}(|\mathbf{x}_i - \mathbf{x}_j|, \nu_i, \nu_j).$$

The Legendre transform of the Lagrangian \mathcal{L} is always well-defined, assuming $\underline{\Omega}(\nu)$ is symmetric and positive definite for all $\nu \in \mathcal{M}$.



We will also denote $\Gamma_i := (\mathbf{x}_i, \underline{p}_i, \nu_i, \underline{s}_i)$ the phase space point of the *i*-th constituent, and introduce

$$\pi\left(\{\mathsf{\Gamma}_i\}_{i=1}^{\mathsf{N}}
ight)\coloneqq\sum_{i=1}^{\mathsf{N}}\delta\left(\mathsf{\Gamma}_i-\mathsf{\Gamma}_i^*(t)
ight)$$

the Klimontovich distribution function, where $\Gamma_i^*(t)$ is the configuration of the *i*-th constituent at time *t*. We will denote π_s the marginals of the Klimontovich distribution function, with respect to $\Gamma^{(s)} = (\Gamma_{s+1}, \dots, \Gamma_N)$, i.e.

$$\pi_{s}\left(\{\mathsf{\Gamma}_{i}\}_{i=1}^{s}\right) \coloneqq \int \pi\left(\{\mathsf{\Gamma}_{i}\}_{i=1}^{\mathsf{N}}\right) d\mathsf{\Gamma}^{(s)}.$$

The distribution function π_s is called the *s*-particle distribution function, and represents the probability of finding *s* particles in the phase space point $\Gamma_1, \ldots, \Gamma_s$.

BOGOLIUBOV-BORN-GREEN-KIRKWOOD-YVON HIERARCHY

An Introduction to the Theory of the Boltzmann Equation, (S. Harris), Statistical Physics of Particles, (M. Kardar), Statistical Mechanics, 2nd Edition (K. Huang).

Let f_s denote the normalised π_s . We obtain the following expression for the BBGKY hierarchy,

$$\begin{aligned} \frac{\partial f_{s}}{\partial t} + \{\pi_{s}, \mathcal{H}_{s}\} &= \int \sum_{i=1}^{s} \frac{\partial f_{s+1}}{\partial \underline{p}_{i}} \cdot \frac{\partial \mathcal{W}(|\mathbf{x}_{i} - \mathbf{x}_{s+1}|, \nu_{i}, \nu_{s+1})}{\partial \mathbf{x}_{i}} d\Gamma_{s+1} \\ &+ \int \sum_{i=1}^{s} \frac{\partial f_{s+1}}{\partial \underline{\zeta}_{i}} \cdot \frac{\partial \mathcal{W}(|\mathbf{x}_{i} - \mathbf{x}_{s+1}|, \nu_{i}, \nu_{s+1})}{\partial \nu_{i}} d\Gamma_{s+1}, \end{aligned}$$

where
$$\mathcal{H}_s = \left(\sum_{i=1}^s \frac{|p_i|^2}{2m} + \frac{1}{2}\underline{\varsigma_i} \cdot \underline{\Omega}(\nu)^{-1}\underline{\varsigma_i}\right) + \sum_{1 \le i < j \le s} \mathcal{W}(|\mathbf{x}_i - \mathbf{x}_j|, \nu_i, \nu_j).$$



BOGOLIUBOV-BORN-GREEN-KIRKWOOD-YVON HIERARCHY

The first two terms of the BBGKY hierarchy, under the assumption that there are no three-body interactions, amount to

$$\begin{aligned} &\frac{\partial f_1}{\partial t} + \frac{\underline{p}_1}{\underline{m}} \cdot \frac{\partial f_1}{\partial \mathbf{x}_1} + \underline{\Omega}(\nu_1)^{-1} \underline{\varsigma}_1 \frac{\partial f_1}{\partial \nu_1} = \\ &+ \int \frac{\partial \mathcal{W}(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2)}{\partial \mathbf{x}_1} \Big(\frac{\partial f_2}{\partial \underline{p}_1} - \frac{\partial f_2}{\partial \underline{p}_2} \Big) \\ &+ \int \frac{\partial \mathcal{W}(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2)}{\partial \nu_1} \Big(\frac{\partial f_2}{\partial \underline{\varsigma}_1} - \frac{\partial f_2}{\partial \underline{\varsigma}_2} \Big) \end{aligned}$$

$$\begin{aligned} \frac{\partial f_2}{\partial t} &+ \frac{\underline{p}_1}{\underline{m}} \cdot \frac{\partial f_2}{\partial \mathbf{x}_1} + \underline{\Omega}(\nu_1)^{-1} \underline{\varsigma}_1 \cdot \frac{\partial f_2}{\partial \nu_1} \\ &+ \frac{\underline{p}_2}{\underline{m}} \cdot \frac{\partial f_2}{\partial \mathbf{x}_2} + \underline{\Omega}(\nu_2)^{-1} \underline{\varsigma}_2 \cdot \frac{\partial f_2}{\partial \nu_2} \\ &- \frac{\partial \mathcal{W}(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2)}{\partial \mathbf{x}_1} \Big(\frac{\partial f_2}{\partial \underline{p}_1} - \frac{\partial f_2}{\partial \underline{p}_2} \Big) \\ &- \frac{\partial \mathcal{W}(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2)}{\partial \nu_1} \Big(\frac{\partial f_2}{\partial \underline{\varsigma}_1} - \frac{\partial f_2}{\partial \underline{\varsigma}_2} \Big) = 0 \end{aligned}$$



BOGOLIUBOV-BORN-GREEN-KIRKWOOD-YVON HIERARCHY

To highlight the same timescale separation in the second term of the hierarchy we introduce fast and slow varying coordinates, i.e.

$${f x} = {f x}_2 - {f x}_1, \qquad {f X} = rac{1}{2} \left({f x}_2 + {f x}_1
ight).$$

We then boxed the terms that are quickly varying in the second equation of the BBGKY hierarchy, i.e.

$$\frac{\partial f_2}{\partial t} + \frac{1}{2} \frac{\underline{p}_2 + \underline{p}_1}{m} \cdot \frac{\partial f_2}{\partial \underline{X}} + \underline{\underline{\Omega}}(\nu_1)^{-1} \underline{\varsigma}_1 \cdot \frac{\partial f_2}{\partial \nu_1} + \underline{\underline{\Omega}}(\nu_2)^{-1} \underline{\varsigma}_2 \cdot \frac{\partial f_2}{\partial \nu_2} + \left| \frac{\underline{p}_2 - \underline{p}_1}{m} \cdot \frac{\partial f_2}{\partial \mathbf{x}} \right| \\ - \left[\frac{\partial \mathcal{W}(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2)}{\partial \mathbf{x}_1} \cdot \left(\frac{\partial f_2}{\partial \underline{p}_1} - \frac{\partial f_2}{\partial \underline{p}_2} \right) \right] - \frac{\partial \mathcal{W}(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2)}{\partial \nu_1} \cdot \frac{\partial f_2}{\partial \underline{\varsigma}_1} = 0$$





Using the embedding results previously discussed, we can use the fast and slow varying coordinates also for the order parameters, i.e.

$$\underline{n} = \underline{\nu}_2 - \underline{\nu}_1, \qquad \underline{N} = \frac{1}{2} \left(\underline{\nu}_2 + \underline{\nu}_1 \right).$$

We then introduce $\underline{A} = \frac{1}{2} \left(\underline{\Omega}_2(\underline{\nu}a_1)^{-1} \underline{\varsigma}_1 + \underline{\Omega}_2(\underline{\nu}_2)^{-1} \underline{\varsigma}_2 \right)$, $\underline{B} = \left(\underline{\Omega}_2(\underline{\nu}_2)^{-1} \underline{\varsigma}_2 - \Omega_1(\underline{\nu}_1)^{-1} \underline{\varsigma}_1 \right)$, i.e.

$$\begin{split} &\frac{\partial f_2}{\partial t} + \frac{1}{2} \frac{\underline{P}_2 + \underline{P}_1}{m} \cdot \frac{\partial f_2}{\partial \underline{X}} + \underline{A} \cdot \frac{\partial f_2}{\partial \underline{N}} + \boxed{\underline{B} \cdot \frac{\partial f_2}{\partial n}} + \underbrace{\underline{\underline{P}_2 - \underline{P}_1}_{m} \cdot \frac{\partial f_2}{\partial \mathbf{x}}}_{m} \\ &- \boxed{\frac{\partial \mathcal{W}(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2)}{\partial \mathbf{x}_1} \cdot \left(\frac{\partial f_2}{\partial \underline{P}_1} - \frac{\partial f_2}{\partial \underline{P}_2}\right)} - \underbrace{\frac{\partial \mathcal{W}(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2)}{\partial \nu_1} \cdot \left(\frac{\partial f_2}{\partial \underline{\zeta}_1} - \frac{\partial f_2}{\partial \underline{\zeta}_2}\right)}_{D} = 0. \end{split}$$



A VLASOV-TYPE EQUATION





From the separation of timescales in the BBGKY hierarchy we obtain the following identity,

$$\frac{\underline{p}_2 - \underline{p}_1}{m} \cdot \frac{\partial f_2}{\partial \mathbf{x}} = \frac{\partial \mathcal{W}}{\partial \mathbf{x}_1} \left(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2 \right) \cdot \left(\frac{\partial f_2}{\partial \underline{p}_1} - \frac{\partial f_2}{\partial \underline{p}_2} \right)$$

Substituting this identity in the second equation of the BBGKY hierarchy we obtain the following equation,

$$\begin{aligned} \frac{\partial f_1}{\partial t} + \frac{\underline{p}_1}{m} \cdot \frac{\partial f_1}{\partial \mathbf{x}_1} + \underline{\underline{\Omega}}(\nu_1)^{-1} \underline{\underline{\zeta}}_1 \cdot \frac{\partial f_1}{\partial \nu_1} &= \int \frac{\underline{p}_2 - \underline{p}_1}{m} \cdot \frac{\partial f_2}{\partial \mathbf{x}} d\Gamma_2 \\ &+ \int \frac{\partial \mathcal{W}(|\mathbf{x}_1 - \mathbf{x}_2|, \nu_1, \nu_2)}{\partial \nu_1} \cdot \frac{\partial f_2}{\partial \underline{\zeta}_1} d\Gamma_2. \end{aligned}$$



We might be tempted to assume interactions are **weak**,

 $f_2(\Gamma_1,\Gamma_2,t)=f_1(\Gamma_1,t)f_1(\Gamma_2,t).$

This leads to equations of a **reversible nature**, compatible with **Loschmidt's paradox**.

Thus, we have no guarantee that the system described thermalises to a Maxwellian distribution.



We might be tempted to assume interactions are **weak**,

 $f_2(\Gamma_1,\Gamma_2,t)=f_1(\Gamma_1,t)f_1(\Gamma_2,t).$

This leads to equations of a **reversible nature**, compatible with **Loschmidt's paradox**.

Thus, we have no guarantee that the system described thermalises to a Maxwellian distribution.

Weak-order Interactions

We will say that a kinetic equation is governed by **weak-order interactions** if the derivative of the two-particle distribution function factorises as,

 $\begin{aligned} \partial_{\nu_i} f_2(\Gamma_1, \Gamma_2, t) &= f_1(\Gamma_i, t) \partial_{\nu_i} f_1(\Gamma_j, t), \\ \partial_{\varsigma_i} f_2(\Gamma_1, \Gamma_2, t) &= f_1(\Gamma_j, t) \partial_{\varsigma_i} f_1(\Gamma_i, t), \end{aligned}$

for $i \neq j$ and i, j = 1, 2.

VLASOV-TYPE EQUATION



Under the assumption of weak-order interactions we can rewrite the first equation of the BBGKY hierarchy as,

$$\frac{\partial f}{\partial t} + \underline{\dot{x}} \cdot \nabla_{\mathbf{x}} f + \underline{\dot{\nu}} \cdot \nabla_{\nu} f + \mathcal{V} \cdot \nabla_{\varsigma} f = C[f, f],$$

where the collision operator C[f, f] can be written using the transition "probability" W as,

$$C[f_{1}, f_{1}] = \int d\Xi'_{1} d\Xi'_{2} d\Xi_{2} \int_{0}^{\frac{\pi}{2}} \int_{0}^{2\pi} W(\Xi'_{1}, \Xi'_{2} \mapsto \Xi_{1}, \Xi_{2}) f_{1}(\Gamma'_{1}, t) f_{1}(\Gamma'_{2}, t) - W(\Xi_{1}, \Xi_{2} \mapsto \Xi'_{1}, \Xi'_{2}) f_{1}(\Gamma_{1}, t) f_{1}(\Gamma_{2}, t) d\theta_{2} d\varphi_{2}$$

BOLTZMANN INEQUALITY AND THERMALISATION



J. Stat. Phys. Volume 26, 795–801 (C. Cercignani, M. Lampis).

As we said before the collision operator C[f, f] considered here guarantees that the system thermalises to a Maxwellian distribution. In particular, we can prove

 $\int d\Xi \log(f(\Gamma, t))C[f, f] \leq 0,$

which is a generalisation of the **Boltzmann inequality** for Boltzmann's equation with internal degrees of freedom. Following the classical calculus of variation approach we can prove that the unique Maxwellian with prescribed collision invariants is

$$\bar{f}(\Gamma, t) = \exp\left(a + \underline{b} \cdot \underline{p} + c(\underline{p} \times \mathbf{x} + \underline{w}_{\nu} \times \underline{\varsigma}) + d(m^{-1}\underline{p} \cdot \underline{p} + \varsigma \cdot \underline{\underline{\Omega}}(\nu)^{-1}\varsigma)\right).$$

SPACE HOMOGENEOUS VLASOV-TYPE EQUATION



We are interested in the time evolution of the distribution $f(\underline{v}, \nu, \underline{\dot{\nu}}, t)$, $v \in \mathbb{R}^2$, $\nu \in \mathcal{M}$, $\underline{\varsigma} \in T_{\nu}\mathcal{M}$, and $t \ge 0$, solution to the space-homogeneous equation

$$\frac{\partial f}{\partial t} + \underline{\underline{\Omega}}(\nu)^{-1} \underline{\underline{\varsigma}} \cdot \nabla_{\nu} f + \mathcal{V} \cdot \nabla_{\underline{\varsigma}} f = \frac{1}{\tau} \mathcal{C}[f, f],$$

where τ has been obtained rescaling the collision frequency, and as collision operator we consider the one associated with Maxwellian molecules, i.e.

$$C[f,f] = \int d_{\underline{\varsigma}_2} d\underline{\nu}_2 d\nu_2 f' f'_* - \int d_{\underline{\varsigma}_2} d\underline{\nu}_2 d\nu_2 ff_*, \qquad (1)$$

complemented with initial conditions $f(\underline{\nu}, \nu, \underline{\varsigma}, 0) = f_0(\underline{\nu}, \nu, \underline{\varsigma})$ and where we will denote $f_* = f(\underline{\nu}_2, \nu_2, \underline{\varsigma}_2, t)$, and f', f'_* are the distributions depending on the post interaction coordinates.

OXFORD Mathematical hstitute

DIRECT SIMULATIONS MONTE CARLO (DSMC)

We consider discretization of the time interval $[0, T_f]$, with $T_f > 0$ final simulation time, of step $\Delta t > 0$ such that $t^n = n\Delta t$. By $f^n(\underline{v}, \nu, \underline{\varsigma})$ we denote an approximation of $f(\underline{v}, \nu, \underline{\varsigma}, t^n)$ at the *n*-th time step and we apply a splitting method between the Vlasov-type transport operator and the collisional operator.

DIRECT SIMULATIONS MONTE CARLO (DSMC)



We consider discretization of the time interval $[0, T_f]$, with $T_f > 0$ final simulation time, of step $\Delta t > 0$ such that $t^n = n\Delta t$. By $f^n(\underline{v}, \nu, \underline{\varsigma})$ we denote an approximation of $f(\underline{v}, \nu, \underline{\varsigma}, t^n)$ at the *n*-th time step and we apply a splitting method between the Vlasov-type transport operator and the collisional operator.

Transport $\mathcal{T}_{\Delta t}(\cdot)$
We solve the Vlasov-type step \hat{f} =
$\mathcal{T}_{\Delta t}(f^n)$
$\begin{cases} \frac{\partial \hat{f}}{\partial t} + \Omega(\nu)^{-1} \underline{\varsigma} \cdot \nabla_{\nu} \hat{f} + \mathcal{V} \cdot \nabla_{\underline{\varsigma}} \hat{f} = 0 \\ \hat{\varsigma} \end{cases}$
$\int \hat{f}(\underline{\nu}, \nu, \underline{\varsigma}, 0) = f^n(\underline{\nu}, \nu, \underline{\varsigma})$

DIRECT SIMULATIONS MONTE CARLO (DSMC)



We consider discretization of the time interval $[0, T_f]$, with $T_f > 0$ final simulation time, of step $\Delta t > 0$ such that $t^n = n\Delta t$. By $f^n(\underline{v}, \nu, \underline{\varsigma})$ we denote an approximation of $f(\underline{v}, \nu, \underline{\varsigma}, t^n)$ at the *n*-th time step and we apply a splitting method between the Vlasov-type transport operator and the collisional operator.

Transport $\mathcal{T}_{\Delta t}(\cdot)$

We solve the Vlasov-type step
$$\hat{f} = \mathcal{T}_{\Delta t}(f^n)$$

$$\begin{cases} \frac{\partial \hat{f}}{\partial t} + \Omega(\nu)^{-1} \underline{\varsigma} \cdot \nabla_{\nu} \hat{f} + \mathcal{V} \cdot \nabla_{\underline{\varsigma}} \hat{f} = 0\\ \hat{f}(\underline{\nu}, \nu, \underline{\varsigma}, 0) = f^n(\underline{\nu}, \nu, \underline{\varsigma}) \end{cases}$$

Collision $\mathcal{Q}_{\Delta t}(\cdot)$

We then solve the collision step $\hat{f} = Q_{\Delta t}(\hat{f})$ with initial data given by the solution of the previous step

$$\begin{cases} \tau \frac{\partial \hat{f}}{\partial t} = \mathcal{C}[\hat{f}, \hat{f}] \\ \hat{f}(\underline{\nu}, \nu, \underline{\varsigma}, \mathbf{0}) = \hat{f}(\underline{\nu}, \nu, \underline{\varsigma}, \Delta t). \end{cases}$$

DIRECT SIMULATIONS MONTE CARLO (DSMC)



We consider discretization of the time interval $[0, T_f]$, with $T_f > 0$ final simulation time, of step $\Delta t > 0$ such that $t^n = n\Delta t$. By $f^n(\underline{v}, \nu, \underline{\varsigma})$ we denote an approximation of $f(\underline{v}, \nu, \underline{\varsigma}, t^n)$ at the *n*-th time step and we apply a splitting method between the Vlasov-type transport operator and the collisional operator.

Transport $\mathcal{T}_{\Delta t}(\cdot)$

We solve the Vlasov-type step
$$\hat{f} = \mathcal{T}_{\Delta t}(f^n)$$

$$\begin{cases} \frac{\partial \hat{f}}{\partial t} + \Omega(\nu)^{-1} \underline{\varsigma} \cdot \nabla_{\nu} \hat{f} + \mathcal{V} \cdot \nabla_{\underline{\varsigma}} \hat{f} = 0\\ \hat{f}(\underline{\nu}, \nu, \underline{\varsigma}, 0) = f^n(\underline{\nu}, \nu, \underline{\varsigma}) \end{cases}$$

Collision $\mathcal{Q}_{\Delta t}(\cdot)$

We then solve the collision step $\hat{f} = Q_{\Delta t}(\hat{f})$ with initial data given by the solution of the previous step

$$\begin{cases} \tau \frac{\partial \hat{f}}{\partial t} = \mathcal{C}[\hat{f}, \hat{f}] \\ \hat{f}(\underline{\nu}, \nu, \underline{\varsigma}, 0) = \hat{f}(\underline{\nu}, \nu, \underline{\varsigma}, \Delta t). \end{cases}$$

The first order in time splitting finally reads $f^{n+1}(\underline{\nu}, \nu, \underline{\varsigma}) = \mathcal{Q}_{\Delta t}(\mathcal{T}_{\Delta t}(f^n)(\underline{\nu}, \nu, \underline{\varsigma})).$



We introduce an approximation of the distribution function with a sample of N particles identified by their velocities $\underline{\nu}_i^n$, order parameter ν_i^n , and conjugate momentum $\underline{\varsigma}_i^n$ at the time t^n , for i = 1, 2, ..., N,

$$f^n(\underline{\nu}, \nu, \underline{\varsigma}) pprox f^{n,N}(\nu, \underline{\varsigma}) = \sum_{i=1}^N \delta(\nu - \nu_i(t^n)) \otimes \delta(\underline{\varsigma} - \underline{\varsigma}_i(t^n)).$$

The Vlasov-type transport step $\mathcal{T}_{\Delta t}(\cdot)$ is solved by considering the characteristic equations associated to the operator, which as discussed in the previous section, result in a system of (time-continuos) ODEs

$$rac{d\underline{
u}_i}{dt} = \underline{\varsigma}_i, \qquad rac{d\underline{\varsigma}_i}{dt} = \mathcal{V}(\nu_i, \underline{\varsigma}_i).$$

This system is solved, at the time discrete level, with a classical first order Euler scheme for the time derivative.



The collisional step $Q_{\Delta t}(\cdot)$ is solved with a classical Nanbu-Babovsky DSMC approach. First, we rewrite the collisional operator to highlight the gain and loss part integrating the second term in (1)

$$G-L=\int d\underline{\varsigma}_2\,d\underline{\nu}_2\,d\nu_2f'f'_*-f,$$

and then we discretize the time derivative with a first order in time Euler scheme to obtain

$$f^{n+1} = \left(1 - \frac{\Delta t}{\tau}\right) f^n + \frac{\Delta t}{\tau} \int d\underline{\varsigma}_2 \, d\underline{\nu}_2 \, d\nu_2 f' f'_*.$$

We have thus rewritten f^{n+1} as a convex combination of f^n and the gain term, i.e. we will consider all the particles in the system with probability $\frac{\Delta t}{\tau}$ we will update the velocity, order parameter and conjugate momentum according to the binary law relating the pre and post interaction velocities, order parameters and conjugate momenta.

Oxford Mathematics



In the context of of rod-like molecules, with vanishing girth, we can explicitly compute the Vlasov-type force \mathcal{V} and the transport term to obtain the following equation

$$rac{\partial f}{\partial t} + \omega
abla_{ heta} f + \mathcal{V} \cdot
abla_{\omega} f = \iiint \left(f' f'_* - f f_*
ight) dv_* d heta_* d\omega_*,$$

where $f = f(\underline{v}, \theta, \omega, t)$, $f_* = f(\underline{v}_*, \theta_*, \omega_*, t)$, and f', f'_* are the distributions depending on the post interaction coordinates given by

$$egin{aligned} & \underline{v}' = \underline{v} - (1+e_v) rac{J}{m} \underline{n}, & \underline{v}'_* = \underline{v}_* + (1+e_v) rac{J}{m} \underline{n}, \\ & \omega' = \omega - (1+e_\omega) J \mathbb{I}^{-1} (r imes \underline{n}), & \omega'_* = \omega_* + (1+e_\omega) J \mathbb{I}^{-1}_* (r_* imes \underline{n}), \end{aligned}$$

with

$$J = -\frac{V \cdot \underline{n}}{\frac{2}{m} + \left[\mathbb{I}^{-1}(r \times \underline{n}) \times r + \mathbb{I}^{-1}_{*}(r_{*} \times \underline{n}) \times r_{*}\right] \cdot \underline{n}}.$$

Notice that $\theta' = \theta$ and $\theta'_* = \theta_*$ since the angles are not changed by the collisional operator.

A Vlasov-type equation

AN EXAMPLE: NEMATIC LIQUID CRYSTALS DSMC SIMULATIONS



Figure: Test 1 - Zero Potential. We here consider the case of no transport, i.e. $\mathcal{V}(\nu,\varsigma) = 0$.



OXFORD Mathematical

AN EXAMPLE: NEMATIC LIQUID CRYSTALS DSMC SIMULATIONS



Figure: Test 1 - Zero Potential. We here consider the case of no transport, i.e. $\mathcal{V}(\nu,\varsigma) = 0$.

Let us consider the mean-field potential is given by

$$\mathcal{W}(\nu,\underline{\varsigma}) = \frac{1}{2} \alpha \left(\underline{\nu} - \underline{\hat{\nu}} \right) \cdot \left(\underline{\nu} - \underline{\hat{\nu}} \right) + \beta \, \underline{\nu} \cdot \underline{\varsigma}.$$

Under this hypothesis the Vlasov-type force can be computed to be

$$\mathcal{V}(\nu,\underline{\varsigma}) = -\alpha \left(\underline{\nu} - \underline{\hat{\nu}}\right) - \beta \underline{\varsigma}.$$

This system of ODEs can be recasted as linear system of ODEs, i.e.

$$\begin{bmatrix} \frac{d\underline{\nu}_i}{dt} \\ \frac{d\underline{\varsigma}_i}{dt} \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -\alpha & -\beta \end{bmatrix} \begin{bmatrix} \underline{\nu}_i \\ \underline{\varsigma}_i \end{bmatrix} + \alpha \begin{bmatrix} 0 \\ \underline{\hat{\nu}} \end{bmatrix}.$$

We can immediately see that the fixed points of the system is unique and it is given by $\underline{\nu} = \hat{\underline{\nu}}$ and $\underline{\varsigma} = 0$. It remains to study the stability of the fixed point, which can be done by studying the eigenvalues of the Jacobian of the system which are given by

$$\lambda_{1,2} = \frac{-\beta \pm \sqrt{\beta^2 - 4\alpha}}{2}$$

A Vlasov-type equation

AN EXAMPLE: NEMATIC LIQUID CRYSTALS DSMC SIMULATIONS



Figure: Test 2 - Linearised Potential. We here consider the case of a linear potential, i.e. $\mathcal{W}(\nu, \underline{\varsigma}) = \alpha (\underline{\nu} - \underline{\hat{\nu}}) \cdot (\underline{\nu} - \underline{\hat{\nu}}) + \beta \underline{\nu} \cdot \underline{\varsigma}$, with $\alpha = 0.1$ and $\beta = 0.1$. We can observe from the right most plot that the system exhibits alignment.



AN EXAMPLE: NEMATIC LIQUID CRYSTALS DSMC SIMULATIONS





Figure: Test 2 - Linearised Potential. We here consider the case of a linear potential, i.e. $\mathcal{W}(\nu, \underline{\varsigma}) = \alpha (\underline{\nu} - \underline{\hat{\nu}}) \cdot (\underline{\nu} - \underline{\hat{\nu}}) + \beta \underline{\nu} \cdot \underline{\varsigma}$, with $\alpha = 0.1$ and $\beta = 0.1$. We can observe from the right most plot that the system exhibits alignment.

A Vlasov-type equation

AN EXAMPLE: NEMATIC LIQUID CRYSTALS DSMC SIMULATIONS



Figure: Test 3 - Non-Linear Potential. We here consider the case of a non-linear potential, i.e. $\mathcal{W}(\theta) = \alpha \cos(\theta - \hat{\theta})$, with $\hat{\theta} = \arctan(\hat{\underline{\nu}}_y, \hat{\underline{\nu}}_x)$, $\alpha = 1$ and $\hat{\underline{\nu}} = \frac{1}{N} \sum_{i=1}^{N} \underline{\nu}_i$. We can observe from the right most plot that the system exhibits alignment.



AN EXAMPLE: NEMATIC LIQUID CRYSTALS DSMC SIMULATIONS





Figure: Test 3 - Non-Linear Potential. We here consider the case of a non-linear potential, i.e. $\mathcal{W}(\theta) = \alpha \cos(\theta - \hat{\theta})$, with $\hat{\theta} = \arctan(\underline{\hat{\nu}}_y, \underline{\hat{\nu}}_x)$, $\alpha = 1$ and $\underline{\hat{\nu}} = \frac{1}{N} \sum_{i=1}^{N} \underline{\nu}_i$. We can observe from the right most plot that the system exhibits alignment.

THANK YOU!

A Kinetic Framework for Fluids with Ordering

Umberto Zerbinati^{*}, joint work with: J. A. Carillo^{*}, P. E. Farrell^{*}, A. Medaglia^{*}.